

mean hitting time*

CWoo[†]

2013-03-21 17:22:12

Let $(X_n)_{n \geq 0}$ be a Markov chain with transition probabilities p_{ij} where i, j are states in an indexing set I . Let H^A be the hitting time of $(X_n)_{n \geq 0}$ for a subset $A \subseteq I$. That is, H^A is the random variable of the time it takes for the chain to first reach a state in A .

Define the *mean hitting time* of A given the chain starts in state i to be

$$k_i^A := E(H^A | X_0 = i).$$

Proposition 1. *The mean hitting times are the minimal non negative solution to:*

$$k_i^A = \begin{cases} 0 & i \in A \\ 1 + \sum_{j \in I} p_{ij} k_j^A & i \notin A \end{cases}$$

Remark. In this case, a solution is minimal if for any non negative solution $\{y_i | i \in I\}$ we have $y_i \geq k_i^A$ for all $i \in I$.

Proof. If $i \in A$, then $H^A = \inf\{n \geq 0 | X_n \in A\} \equiv 0$, which means $k_i^A = 0$ (the chain is certain to be in a state in A at step $n = 0$).

If $i \notin A$ we condition on the first step:

$$\begin{aligned} k_i^A &= E(H^A | X_0 = i) \\ &= \sum_{j \in I} P(X_1 = j | X_0 = i) E(H^A | X_0 = i, X_1 = j) \\ &= \sum_{j \in I} p_{ij} E(H^A | X_1 = j) \quad (\text{by the Markov property}) \\ &= \sum_{j \in I} p_{ij} (1 + k_j^A) \\ &= 1 + \sum_{j \in I} p_{ij} k_j^A \end{aligned}$$

* $\langle \text{MeanHittingTime} \rangle$ created: $\langle 2013-03-21 \rangle$ by: $\langle \text{CWoo} \rangle$ version: $\langle 35807 \rangle$ Privacy setting: $\langle 1 \rangle$ $\langle \text{Theorem} \rangle$ $\langle 60J10 \rangle$

[†]This text is available under the Creative Commons Attribution/Share-Alike License 3.0. You can reuse this document or portions thereof only if you do so under terms that are compatible with the CC-BY-SA license.

So the k_i^A satisfy the given equations.

Now suppose that $\{y_i | y \in I\}$ is any non-negative solution to the equations. Then for $i \in A$ we have $k_i^A = y_i = 0$. If $i \notin A$, then

$$\begin{aligned}
y_i &= 1 + \sum_{j \in I} p_{ij} y_j \\
&= 1 + \sum_{j \notin A} p_{ij} (1 + \sum_{k \notin A} p_{jk} y_k) \\
&= 1 + \sum_{j \notin A} p_{ij} + \sum_{j \notin A} \sum_{k \notin A} p_{ij} p_{jk} y_k \\
&= 1 + q_1 + q_2 + \cdots + q_n + \sum \cdots \sum p_{ij} \cdots p_{uv} y_v,
\end{aligned}$$

where $q_n = P(X_1 \notin A, X_2 \notin A, \dots, X_n \notin A | X_0 = i)$ is the probability that the chain X does not enter A in the first n steps after the initial state i .

y_i is non negative by assumption, therefore so is the final term, and so

$$y_i \geq 1 + q_1 + q_2 + \cdots + q_n.$$

Since n is arbitrary, by taking the limit $n \rightarrow \infty$, we have that

$$y_i \geq \lim_{n \rightarrow \infty} (1 + q_1 + q_2 + \cdots + q_n) \geq k_i^A.$$

So $y_i \geq k_i^A$ for all $i \in I$ and therefore $\{k_i^A | i \in I\}$ is the minimal solution. \square