

preadditive category*

CWoo†

2013-03-21 20:31:40

0.1 Ab-Category

A category \mathcal{C} is an *ab-category* or *ab-category* if

1. for every pair of objects A, B of \mathcal{C} , there is a binary operation called addition, written $+_{(A,B)}$ or simply $+$, defined on $\text{hom}(A, B)$,
2. the set $\text{hom}(A, B)$, together with $+$ is an abelian group,
3. (left distributivity) if $f, g \in \text{hom}(A, B)$ and $h \in \text{hom}(B, C)$, then $h(f + g) = hf + hg$,
4. (right distributivity) if $f, g \in \text{hom}(A, B)$ and $h \in \text{hom}(C, A)$, then $(f + g)h = fh + gh$.

In a nutshell, an *ab-category* is a category in which every hom set in \mathcal{C} is an abelian group such that morphism composition distributes over addition. Ab in the name stands for abelian, clearly indicative of the second condition above.

Since a group has a multiplicative (or additive if abelian) identity, $\text{hom}(A, B) \neq \emptyset$ for every pair of objects A, B in \mathcal{C} . Furthermore, each $\text{hom}(A, B)$ contains a unique morphism, written $0_{(A,B)}$, as the additive identity of $\text{hom}(A, B)$. Because the subset

$$\{f \cdot 0_{(A,B)} \mid f \in \text{hom}(B, C)\}$$

of $\text{hom}(A, C)$ is also a subgroup by right distributivity, and the additive identity of a subgroup coincides with the additive identity of the group, we have the following identity

$$0_{(B,C)}0_{(A,B)} = 0_{(A,C)}.$$

There are many examples of ab-categories, including the category of abelian groups, the category of R -modules (R a ring), the category of chain complexes, and the category of rings (not necessarily containing a multiplicative identity).

**(PreadditiveCategory)* created: *(2013-03-21)* by: *(CWoo)* version: *(37913)* Privacy setting: *(1)* *(Definition)* *(18E05)*

†This text is available under the Creative Commons Attribution/Share-Alike License 3.0. You can reuse this document or portions thereof only if you do so under terms that are compatible with the CC-BY-SA license.

However, the category of rings with 1 is not an ab-category (see below for more detail). Nevertheless, a unital ring R itself considered as a category is an ab-category, as the ring of endomorphisms clearly forms an abelian group. It is in fact a ring! This can be seen as a special case of the fact that, in an ab-category, $\text{End}(A) = \text{hom}(A, A)$ is always a ring (with 1). So, conversely, an ab-category with one object is a ring with 1, whose morphisms are elements of the ring.

0.2 Preadditive Category

If an ab-category has an initial object, that object is also a terminal object. By duality, the converse is also true. Therefore, in an ab-category, initial object, terminal object, and zero object are synonymous. In the category \mathcal{R} of unital rings, \mathbb{Z} is an initial object, but it has no terminal object, therefore \mathcal{R} is not an ab-category.

An ab-category with a zero object O is called a *preadditive category*.

In a preadditive category, the groups $\text{hom}(A, O)$ and $\text{hom}(O, B)$ are trivial groups by the definition of the zero object O . Therefore, the zero morphism in $\text{hom}(A, B)$ is also the additive identity of $\text{hom}(A, B)$:

$$0_{(A,B)} = 0_{(O,B)}0_{(A,O)} = A \longrightarrow O \longrightarrow B.$$

Most of the examples of ab-categories are readily seen to be preadditive. If a preadditive category R has only one object, we see from above that it must be a ring. But this object must also be a zero object, so that $\text{End}(R)$ must be trivial, which means R itself must be trivial too, $R = 0$!

Remark. In some literature, a preadditive category is an ab-category, and some do not insist that a preadditive category contains a zero object. Here, we choose to differentiate the two.