

Boolean quotient algebra*

CWoo[†]

2013-03-22 0:51:58

Quotient Algebras via Congruences

Let A be a Boolean algebra. A congruence on A is an equivalence relation Q on A such that Q respects the Boolean operations:

- if aQb and cQd , then $(a \vee c)Q(b \vee d)$
- if aQb , then $a'Qb'$

By de Morgan's laws, we also have aQb and cQd implying $(a \wedge c)Q(b \wedge d)$.

When a is congruent to b , we usually write $a \equiv b \pmod{Q}$.

Let B be the set of congruence classes: $B = A/Q$, and write $[a]Q$, or simply $[a]$ for the congruence class containing the element $a \in A$. Define on B the following operations:

- $[a] \vee [b] := [a \vee b]$
- $[a]' := [a']$

Because Q respects join and complementation, it is clear that these are well-defined operations on B . Furthermore, we may define $[a] \wedge [b] := ([a]' \vee [b]')' = ([a]' \vee [b]')' = [a' \vee b']' = [(a' \vee b)'] = [a \wedge b]$. It is also easy to see that $[1]$ and $[0]$ are the top and bottom elements of B . Finally, it is straightforward to verify that B is a Boolean algebra. The algebra B is called the *Boolean quotient algebra* of A via the congruence Q .

Quotient Algebras via Ideals and Filters

It is also possible to define quotient algebras via Boolean ideals and Boolean filters. Let A be a Boolean algebra and I an ideal of A . Define binary relation \sim on A as follows:

$$a \sim b \quad \text{if and only if} \quad a \Delta b \in I,$$

**BooleanQuotientAlgebra* created: $\langle 2013-03-2 \rangle$ by: $\langle CWoo \rangle$ version: $\langle 40496 \rangle$ Privacy setting: $\langle 1 \rangle$ *Definition* $\langle 06E05 \rangle$ $\langle 03G05 \rangle$ $\langle 06B20 \rangle$ $\langle 03G10 \rangle$

[†]This text is available under the Creative Commons Attribution/Share-Alike License 3.0. You can reuse this document or portions thereof only if you do so under terms that are compatible with the CC-BY-SA license.

where Δ is the symmetric difference operator on A . Then

1. \sim is an equivalence on A , because
 - $a\Delta a = 0 \in I$, so \sim is reflexive
 - $b\Delta a = a\Delta b$, so \sim is symmetric, and
 - if $a \sim b$ and $b \sim c$, then $a \sim c$; to see this, note that $(a-b)\vee(b-c) = ((a-b)\vee b)\wedge((a-b)\vee c') = (a\vee b)\wedge((a-b)\vee c')$. Since the LHS (and hence the RHS) is in I , and that $a \leq a\vee b$ and $c' \leq (a-b)\vee c'$, RHS $\geq a\wedge c' = a-c \in I$ too. Similarly $c-a \in I$ so that $a \sim c$.
2. \sim respects \vee and $'$, because
 - if $a \sim b$ and $c \sim d$, then $(a\vee c) - (b\vee d) = (a\vee c)\wedge(b\vee d)' = (a\vee c)\wedge(b'\wedge d') = (a\wedge(b'\wedge d'))\vee(c\wedge(b'\wedge d')) \leq (a\wedge b')\vee(c\wedge d') \in I$, so that $(a\vee c) - (b\vee d) \in I$ as well. That $(b\vee d) - (a\vee c) \in I$ is proved similarly. Hence $(a\vee c) \sim (b\vee d)$.
 - $a'\Delta b' = a\Delta b$, so \sim preserves $'$.

Thus, \sim is a congruence on A . The quotient algebra A/\sim is called the *quotient algebra* of A via the ideal I , and is often denoted by A/I .

From this congruence \sim , one can re-capture the ideal: $I = [0]$.

Dually, one can obtain a quotient algebra from a Boolean filter. Specifically, if F is a filter of a Boolean algebra A , define \sim on A as follows:

$$a \sim b \quad \text{if and only if} \quad a \leftrightarrow b \in F,$$

where \leftrightarrow is the biconditional operator on A . Then it is easy to show that \sim too is a congruence on A , so that one forms the *quotient algebra* of A via the filter F , denoted by A/F . Of course, an easier approach to this is to realize that F is a filter of A iff $F' := \{a' \mid a \in F\}$ is an ideal of A , and the process of forming A/F' turns out to be identical to A/F .

From \sim , the filter F can be recovered: $F = [1]$.

In fact, given a congruence Q , the congruence class $[0]Q$ is a Boolean ideal and the congruence class $[1]Q$ is a Boolean filter, and that the quotient algebras derived from Q , $[0]Q$ and $[1]Q$ are all the same:

$$A/Q = A/[0]Q = A/[1]Q.$$